



Cytological Distinctions Between *Timun Suri* and Cucumber Discovered by Fluorescence In Situ Hybridization (FISH) Using 45S Ribosomal DNA Gene

Agus Budi Setiawan^{1*)}, Aziz Purwanto¹⁾ and Ari Wibowo²⁾

¹⁾ Genetic and Plant Breeding Lab, Department of Agronomy, Faculty of Agriculture, Universitas Gadjah Mada, Yogyakarta, Indonesia

²⁾ Indonesian Coffee and Cacao Research Institute, Jember, East Java, Indonesia

ARTICLE INFO

Keywords:

45S rDNA

Cucumber

Cucumis melo var. *momordica*

FISH

Timun suri

Article History:

Received: January 25, 2019

Accepted: October 17, 2020

*) Corresponding author:

E-mail: setiawanab@ugm.ac.id

ABSTRACT

The genus *Cucumis* including *timun suri*, melon and cucumber, is an important horticultural crop of flowering plants. The dispute of *timun suri* terminology is leading to false-positive results in the nomenclature of *timun suri* in Indonesia. Although molecular research on these species has been widely conducted, detailed information and precise evidence based on the molecular cytogenetic approach are poorly investigated. The FISH technique was applied to reveal the cytological distinctions among these species. Here, this research conducted physical mapping of 45S ribosomal DNA (rDNA) on six accessions of *Cucumis*. The chromosome number of cucumber is 14 chromosomes, while *timun suri* and melon are 24 chromosomes. The number of 45S rDNA loci in cucumber, melon and *timun suri* was highly conserved. All cucumber accessions in this study had nine to ten 45S rDNA loci with strong and weak signal intensities located at proximal regions of the short arms. In melon and *timun suri* the signals of two pairs of 45S rDNA were located at the terminal and interstitial regions of the short arms. These were clear shreds of evidence to confirm that *timun suri* was cytogenetically closed to melon instead of cucumber. It confirmed that *timun suri* cannot be classified as a cucumber.

INTRODUCTION

The genus *Cucumis* is one of the valuable genera in the *Cucurbitaceae* family. It includes cucumber and melon, two of the most economically valuable and widely cultivated crops in the world (Pitrat, Chauvet, & Foury, 1999). Cucumber (*Cucumis sativus* L.) is a diploid species having 14 chromosomes and well-known as *mentimun* or *timun* in Indonesia (Han et al., 2008). *Timun suri* (local name - Indonesia), based on its terminology, could be classified as a cucumber. However, that reason cannot be scientifically justified. *Timun suri* was closely related to melon (*Cucumis melo* L.) depending on its morphology, isozyme, and RAPD markers rather than to cucumber (Huda & Daryono, 2013; Rahayu & Hartana, 2002).

In addition, the fruit morphology of *timun suri* in Indonesia was closely related to *Cucumis*

melo subsp *agrestis* var. *momordica* same as that has been reported by Manohar & Murthy (2012). Furthermore, the chromosome number of *C. melo* subsp *agrestis* var. *momordica* has been confirmed 24 chromosomes (Setiawan, 2018). Despite those results, the morphology, isozyme, and RAPD have some limitations, particularly morphological traits and isozyme which were affected by the environmental factor. In addition, poor reproducibility of RAPD has been informed by Penner et al. (1993) and Skroch & Nienhuis (1995).

In order to get more insights and evidence to precisely distinguish between *timun suri*, cucumber, and melon, this research has physically mapped the 45S rDNA gene using FISH and compare its chromosomal distribution between those accessions. Physical mapping of 45S rDNA provides remarkable information on the genetic relationships among the

ISSN: 0126-0537 Accredited First Grade by Ministry of Research, Technology and Higher Education of The Republic of Indonesia, Decree No: 30/E/KPT/2018

Cite this as: Setiawan, A. B., Purwanto, A., & Wibowo, A. (2020). Cytological distinctions between *timun suri* and cucumber discovered by fluorescence in situ hybridization (FISH) using 45S ribosomal DNA gene. *AGRIVITA Journal of Agricultural Science*, 42(3), 584–592. <https://doi.org/10.17503/agrivita.v42i3.2142>

Agus Budi Setiawan et al.: Cytogenetic Evidences *Timun Suri* and *Cucumber*.....

closely related species in several genera such as *Fabaceae*, *Poaceae*, *Solanaceae*, and *Cucumis* (Acosta, Moscone, & Cocucci, 2016; Amosova et al., 2017; Li et al., 2016; She et al., 2015; Zhang et al., 2016). 45S rDNA gene is a tandem repeat sequence and consisting of the 18S, 5.8S, and 25S ribosomal RNA (rRNA) (Ganal & Hemleben, 1986). This rRNA, together with several classes of the protein complex, established the translation factory of protein synthesis (Neves et al., 2005).

The number of rDNA loci and their location in chromosomes have great variations in *Cucumis* species, particularly between melon and cucumber which is clearly distinguishable and can be utilized to clarify and discriminate the cytological marker of these two species in addition to chromosome number (Setiawan, 2018). The number of 45S rDNA loci in melon are conserved two loci and located at Nucleolar Organizing Region (NOR) either in the terminal or interstitial region of the short arms (Li et al., 2016; Liu et al., 2010; Setiawan, 2018; Zhang et al., 2015; Zhang et al., 2016), whereas in cucumber, they were varied from eight to ten loci depending on cultivars (Chen, Staub, Adelberg, & Jiang, 1999; Koo et al., 2010; Zhao et al., 2011).

FISH can be used to check the location and abundance of a DNA sequence through the hybridization of a labeled DNA probe to DNA template in chromosomes (Schwarzacher & Heslop-Harrison, 2000). It can be applied for the identification of chromosomes homologous for karyotyping and phylogenetic analysis, direct physical mapping of sequences on the chromosome, and for studying the recent evolutionary process in the genome (Jiang &

Gill, 2006; Kato, Lamb, & Birchler, 2004). Therefore, FISH is a useful technique to determine the position of 45S rDNA in chromosomes and to compare its chromosomal distribution in *timun suri*, cucumber, and melon. This research provides precise evidence based on the molecular cytogenetic approach to distinguish between *timun suri* and cucumber. It is clear that *timun suri* is classified as melon with nomenclature name *Cucumis melo* subsp. *agrestis* var. *momordica* and has four 45S rDNA loci on somatic metaphase chromosomes.

Molecular research on *timun suri* has been widely conducted, but detailed information and precise evidence based on the molecular cytogenetic approach are poorly investigated. The aim of this research was to evaluate the cytogenetic relativity of *timun suri* and cucumber using the 45S ribosomal DNA gene.

MATERIALS AND METHODS

Plant Materials

One melon accession (NI19), two cucumber landraces including Okute Aodai and RAR 930024, and two accessions of *timun suri* i.e., US 205 and IPB TS1 were used for cytological studies. The seed materials were provided by Dr. Kenji Kato (Okayama University), The National Agriculture and Food Research Organization (NARO) Gene Bank of Japan, and Vegetable Crop Research Institute (Balitsa, Indonesia). (Table 1). The seeds were germinated and maintained at 25°C in the growth chamber. The research was conducted at the laboratory of genetic and plant breeding, Chiba University from March until September 2018.

Table 1. The materials used in this study and their chromosomal features

Cultivar	Accession No	Species	Origin	Status	Chro. Number (2n)	Number of 45S rDNA loci		45S rDNA location
						Strong	Weak	
Litsa Hijau	-	<i>Cucumis sativus</i> L.	IDN	Open pollinated	14	7	3	Prox (SA)
Okute Aodai	32268	<i>Cucumis sativus</i> L.	JPN	Landrace	14	6	3	Prox (SA)
RAR 930024	82368	<i>Cucumis sativus</i> L.	USSR	Landrace	14	6	4	Prox (SA)
NI19	940281	<i>Cucumis melo</i> L.	TCD	Landrace	24	2	2	Ter & Inter (SA)
US 205	PI 182952	<i>Cucumis melo</i> L. Ssp. <i>agrestis</i> var <i>Momordica</i>	IND	Landrace	24	2	2	Ter & Inter (SA)
IPB TS1	-	<i>Cucumis melo</i> L. Ssp. <i>agrestis</i> var <i>Momordica</i>	IDN	Open pollinated	24	4	0	Ter & Inter (SA)

Remarks: Chro. = chromosome; Prox = Proximal region; Ter = Terminal region; Inter = Interstitial region; SA = Short arms; IDN: Indonesia; JPN: Japan; USSR: Uzbekistan, Kazakhstan, and Kyrgyzstan area; TCD: Chad; IND; India

Chromosome Preparations

Timun suri, cucumber, and melon seeds were germinated on filter paper using distilled water, kept in petri dishes in a growth chamber at 25°C. The main root tips were cut (0.5-1 cm). The mitotic and meiotic chromosomes were prepared in accordance with Setiawan, Teo, Kikuchi, Sassa, & Koba (2018) protocol. Modified Carnoy's solution II was used for the mitotic and meiotic chromosomes fixation method. In brief, both of flower buds and root tips were pretreated with 6:3:1 (v/v) ethanol: acetic acid: chloroform for 3-4 hours at room temperature and transferred into 3:1 (v/v) ethanol: acetic acid solution at 4°C for 5 days. The samples were washed for 10 minutes in 1 ml of enzyme buffer (40 ml of 100 mM citric acid + 60 ml of 100 mM sodium citrate, pH 4.8). The anthers were dissected from flower buds under a stereomicroscope using the forceps and the root tips were cut with a razor blade. Both of them were macerated in 15 µl of the enzyme cocktail containing 1% Pectolyase Y-23 (Kyowa Chemical, Osaka, Japan), 2% Pectinase (Sigma), and 4% Cellulose Onozuka RS (Yakult) at 37°C for 1 hour. The enzymes surrounding the roots and anthers were cleaned off using Kimwipe tissue. The squeezing and air-dry method was used for mitotic and meiotic chromosome preparations, respectively, as same as described in Setiawan, Teo, Kikuchi, Sassa, & Koba (2018).

The Probes and FISH Procedure

45S rDNA sequence (pTa71) of wheat was used as the probes (Gerlach & Bedbrook, 1979). The 45S rDNA was cloned in pGEM-T-Easy Vector (Promega), and the plasmid DNA was extracted using HiYield™ Plasmid Kit (RBC Bioscience) in accordance with the manufacturer's instructions. Therefore, the dig-nick translation mix (Roche) was used for labeling the 45S rDNA. The FISH analysis was performed as same as described in Setiawan, Teo, Kikuchi, Sassa, & Koba (2018) method, consisting of pretreatment using RNase A (Qiagen) and pepsin (Sigma) for chromosome slides, 1% paraformaldehyde for refixation, hybridization process including denaturation at 80°C for 2 minutes, and probe detection using anti-digoxigenin rhodamine (Roche).

Image Analysis

The chromosome slides were stained with 4,6-diamidino-2-phenylindole (DAPI) mounted in a

VectaShield antifade solution (Vector Laboratories). The observation was conducted using a fluorescence microscope (Olympus BX53) equipped with a cooled CCD camera (Photometrics CoolSNAP MYO). The processing and editing of FISH images were conducted using Metamorph, Metavue imaging series version 7.8, and Adobe Photoshop CS 6, respectively.

RESULTS AND DISCUSSION

The Conservation of 45S rDNA in *Cucumber*, *Melon*, and *Timun Suri*

The 45S rDNA probe was hybridized on the metaphase chromosomes of six *Cucumis* accessions. All cucumber accessions had 14 chromosomes (Fig. 1a, 1d, 1g). Three accessions of cucumber had variation in 45S rDNA loci consisting of 9 to 10 loci with different signal intensities. Seven strong and three weak signals of 45S rDNA were detected in Litsa Hijau (Fig. 1b-c). Three pairs of strong and three weak signals of 45S rDNA were discovered in Okute Aodai (Fig. 1e-f). While RAR 930024 had three pairs of strong and two pairs of weak signals (Fig. 1h-i), and some of the cucumber cultivars had been reported to have 8 rDNA loci (Chen, Staub, Adelberg, & Jiang, 1999; Lou *et al.*, 2014; Setiawan, 2018).

The presence of eight to ten 45S rDNA loci in wild and domesticated cucumber accessions from different locations implies that the amplification of 45S rDNA loci was occurred before domestication. The different number of 45S rDNA loci in *Phaseolus vulgaris* accessions or in the same genera of *Passiflora*, *Sellaginella*, and *Paspalum* has been reported to have variation in 45S rDNA loci (3 to 9 loci) (De Melo & Guerra, 2003; Marcon, Leão Barros, & Guerra, 2005; Pedrosa-Harand *et al.*, 2006; Vaio, Speranza, Valls, Guerra, & Mazzella, 2005). There are three possible mechanisms of 45S rDNA variation: (1) alteration of chromosome morphology caused by the amplification of satellite DNA without changing the order of other sequences Navrátilová, Neumann, & Macas (2003); (2) translocation, a part of chromosomal rearrangement, have been reported to shift the position of rDNA loci in *Lotus japonicas* (Hayashi *et al.*, 2001); (3) dispersion-amplification-deletion model is proposed to be the responsible for the variation of 45S rDNA loci (Dubcovsky & Dvorák, 1995).




Fig. 1. Physical mapping of 45S rDNA in cucumber, melon, and *timun suri* accessions. Litsa Hijau (a-c); Okute Aodai (d-f); RAR 930024 (g-i); melon NI19 (j-l); *timun suri* US 205 (m-o); *timun suri* IPB TS1 (p-r). 45S rDNA labeled with digoxigenin (green). Weak signals of 45S rDNA (arrowheads). Scale bars = 10 μ m

Although, how the dispersion occurred has not been clear yet, but this mechanism can explain the change in the number of rDNA loci and the dispersion of other repetitive DNA sequences in a genome. Several studies reported that the movement of 45S rDNA is controlled by mobile elements (Raskina, Belyayev, & Nevo, 2004a; 2004b). A recent study has revealed that the LINE element, a non-long terminal repeat (LTR) retrotransposon, has been inserted into 45S rDNA arrays in cucumber accessions, and those accessions had eight to ten of 45S rDNA loci (Setiawan, 2018). Therefore, retrotransposons are thought to cause the variation of 45S rDNA loci in cucumber.

NI19, a melon accession, showed 24 chromosomes with a pair of strong and weak 45S rDNA signals (Fig. 1j-l). US 205 and IPB TS1, two *timun suri* accessions derived from India and Indonesia, respectively, had the same chromosome number (2n = 24) as well as a pair of strong and weak

45S rDNA signals (Fig. 1m-r), and the interphase cells showed that US 205 had two pairs of 45S rDNA signals (Fig. 2a-c). These results are similar as has been reported by (Li et al., 2016; Liu et al., 2010; Zhang et al., 2016). In addition, all *Cucumis* accessions showed different signal intensities of 45S rDNA. The strong and weak of 45S rDNA signals are related to their copy number (Li, Fu, Hu, Huang, & Song, 2006). Gong et al. (2013) reported that weak signals in *Jatropha curcas* indicate the presence of transcriptionally inactive 45S rDNA. Therefore, cucumber has more active 45S rDNA than melon and *timun suri*. The results provide clear evidence that *timun suri* is completely had cytological distinctions in comparison with cucumber. Therefore, a word of *timun* in *timun suri* cannot be justified and classified as cucumber. In fact, *timun suri* is cytogenetically close to melon, and it was classified as *Cucumis melo* L. ssp. *agrestis* var. *momordica*.




Fig. 2. Interphase and pachytene chromosomes of *timun suri* US 205. Physical mapping of 45S rDNA in meiotic Interphase (a-c) and pachytene (d-f) chromosomes. 45S rDNA labeled with digoxigenin (green). Scale bars = 10 μm

Chromosomal Distribution of 45S rDNA in *Timun Suri*, Cucumber, and Melon

FISH results on cucumber metaphase chromosomes in Litsa Hijau, Okute Aodai, and RAR 930024 showed that all 45S rDNA signals including strong and weak signals were located at proximal regions (Fig. 1c, 1f, 1i). These were coincidence with previously reported results (Han et al., 2008; Koo et al., 2010; Lou et al., 2014). Whereas in NI19, US 205, and IPB TS1, the strong and weak signals of 45S rDNA were located at terminal and Interstitial regions of the short arms (Fig. 1l, 1o, 1r), respectively. These locations of 45S rDNA are similar to that have been reported by Zhang et al. (2016).

Higher-resolution FISH of 45S rDNA was detected in the pachytene chromosome (Fig. 2d-f), and it was confirmed that *timun suri* (US 205) had one pair of 45S rDNA in pachytene cell. The homologous recombination constraint may be the cause of the location of 45S rDNA at the terminal region in melon. Therefore, the allelic and non-allelic homologous recombination may cause an important role in the homogenization of intralocus and interloci of 45S rDNA repeat (Roa & Guerra, 2012; Wendel, Schnabel, & Seelanan, 1995). Various location of 45S rDNA was not only found in plant but also in animal and this variation had no detrimental effects for the meiotic segregation and gene stability (Cabrero & Camacho, 2008; Hanson et al., 1996; Pedrosa-Harand et al., 2006).

Timun Suri is Cytogenetically Close to Melon

It is clear that *timun suri* and cucumber have different cytogenetical traits including chromosome number and 45S rDNA loci (Fig. 1 and Table 1). *Timun suri* is classified as a melon group with different nomenclature i.e. *Cucumis melo* ssp. *agrestis* var. *momordica*, while cucumber is closely related to *Cucumis hystrix* L. (Sebastian, Schaefer, Telford, & Renner, 2010). Although the chromosome number between cucumber and *C. hystrix* is different, a fertile amphidiploid from interspecific hybridization, *C. hystivus* ($2n = 4x = 38$), between cucumber and *C. hystrix* has been successfully developed (Chen et al., 2003; Chen et al., 2004; Chen, Staub, Adelberg, Lewis, & Kunkle, 2002; Chen, Staub, Tashiro, Isshiki, & Miyazaki, 1997). Since *timun suri* is closely related to melon, it is possible to conduct the breeding for fruit quality in *timun suri*. Ripe fruit of *timun suri* generally has

the major problem of fruit cracking that can cause significant loss of marketable revenue and yield (Singh et al., 2015). In addition, the sugar content or sweetness is lower than melon (Dhillon et al., 2007). Therefore, it is necessary to conduct the breeding program for *timun suri* to increase its fruit quality and marketable in the future.

CONCLUSION

Timun suri and cucumber are two different species that have significantly cytogenetic differences between them. Physical mapping of 45S rDNA in *timun suri*, cucumber, and melon successfully discovered the fundamental evidence to distinguish among these accessions. *Timun suri* breeding program for fruit quality may be improved due to the close cytogenetic traits between melon and *timun suri*.

ACKNOWLEDGEMENT

We would like to thank to Prof. Kenji Kato (Okayama University, Japan), The National Agriculture and Food Research Organization (NARO) Gene Bank of Japan, and Vegetable Crop Research Institute (Balitsa, Indonesia) for providing us the seed materials.

REFERENCES

- Acosta, M. C., Moscone, E. A., & Cocucci, A. A. (2016). Using chromosomal data in the phylogenetic and molecular dating framework: Karyotype evolution and diversification in *Nierembergia* (Solanaceae) influenced by historical changes in sea level. *Plant Biology*, 18(3), 514–526. <https://doi.org/10.1111/plb.12430>
- Amosova, A. V., Bolsheva, N. L., Zoshchuk, S. A., Twardovska, M. O., Yurkevich, O. Y., Andreev, I. O., ... Muravenko, O. V. (2017). Comparative molecular cytogenetic characterization of seven *Deschampsia* (Poaceae) species. *PLoS ONE*, 12(4), e0175760. <https://doi.org/10.1371/journal.pone.0175760>
- Cabrero, J., & Camacho, J. P. M. (2008). Location and expression of ribosomal RNA genes in grasshoppers: Abundance of silent and cryptic loci. *Chromosome Research*, 16, 595–607. <https://doi.org/10.1007/s10577-008-1214-x>
- Chen, J.-F., Luo, X.-D., Qian, C.-T., Jahn, M. M., Staub, J. E., Zhuang, F.-Y., ... Ren, G. (2004). *Cucumis* monosomic alien addition lines: Morphological,

- cytological, and genotypic analyses. *Theoretical and Applied Genetics*, 108, 1343–1348. <https://doi.org/10.1007/s00122-003-1546-z>
- Chen, J.-F., Staub, J. E., Adelberg, J. W., & Jiang, J. (1999). Physical mapping of 45S rRNA genes in *Cucumis* species by fluorescence *in situ* hybridization. *Canadian Journal of Botany*, 77(3), 389–393. <https://doi.org/10.1139/b98-226>
- Chen, J.-F., Staub, J. E., Qian, C., Jiang, J., Luo, X., & Zhuang, F. (2003). Reproduction and cytogenetic characterization of interspecific hybrids derived from *Cucumis hystrix* Chakr. x *Cucumis sativus* L. *Theoretical and Applied Genetics*, 106, 688–695. <https://doi.org/10.1007/s00122-002-1118-7>
- Chen, J.-F., Staub, J. E., Tashiro, Y., Isshiki, S., & Miyazaki, S. (1997). Successful interspecific hybridization between *Cucumis sativus* L. and *C. hystrix* Chakr. *Euphytica*, 96, 413–419. <https://doi.org/10.1023/A:1003017702385>
- Chen, J.-F., Staub, J., Adelberg, J., Lewis, S., & Kunkle, B. (2002). Synthesis and preliminary characterization of a new species (amphidiploid) in *Cucumis*. *Euphytica*, 123, 315–322. <https://doi.org/10.1023/A:1015095430624>
- De Melo, N. F., & Guerra, M. (2003). Variability of the 5S and 45S rDNA sites in *Passiflora* L. species with distinct base chromosome numbers. *Annals of Botany*, 92(2), 309–316. <https://doi.org/10.1093/aob/mcg138>
- Dhillon, N. P. S., Ranjana, R., Singh, K., Eduardo, I., Monforte, A. J., Pitrat, M., ... Singh, P. P. (2007). Diversity among landraces of Indian snapmelon (*Cucumis melo* var. *momordica*). *Genetic Resources and Crop Evolution*, 54, 1267–1283. <https://doi.org/10.1007/s10722-006-9108-2>
- Dubcovsky, J., & Dvorák, J. (1995). Ribosomal RNA multigene loci: Nomads of the Triticeae genomes. *Genetics*, 140(4), 1367–1377. Retrieved from <https://www.genetics.org/content/140/4/1367>
- Ganal, M., & Hemleben, V. (1986). Comparison of the ribosomal RNA genes in four closely related *Cucurbitaceae*. *Plant Systematics and Evolution*, 154(1), 63–77. <https://doi.org/10.1007/BF00984868>
- Gerlach, W. L., & Bedbrook, J. R. (1979). Cloning and characterization of ribosomal RNA genes from wheat and barley. *Nucleic Acids Research*, 7(7), 1869–1885. <https://doi.org/10.1093/nar/7.7.1869>
- Gong, Z., Xue, C., Zhang, M., Guo, R., Zhou, Y., & Shi, G. (2013). Physical localization and DNA methylation of 45S rRNA gene loci in *Jatropha curcas* L. *PLoS ONE*, 8(12), e84284. <https://doi.org/10.1371/journal.pone.0084284>
- Han, Y.-H., Zhang, Z.-H., Liu, J.-H., Lu, J.-Y., Huang, S.-W., & Jin, W.-W. (2008). Distribution of the tandem repeat sequences and karyotyping in cucumber (*Cucumis sativus* L.) by fluorescence *in situ* hybridization. *Cytogenetic and Genome Research*, 122, 80–88. <https://doi.org/10.1159/000151320>
- Hanson, R. E., Islam-Faridi, M. N., Percival, E. A., Crane, C. F., Ji, Y., McKnight, T. D., ... Price, H. J. (1996). Distribution of 5S and 18S-28S rDNA loci in a tetraploid cotton (*Gossypium hirsutum* L.) and its putative diploid ancestors. *Chromosoma*, 105, 55–61. <https://doi.org/10.1007/BF02510039>
- Hayashi, M., Miyahara, A., Sato, S., Kato, T., Yoshikawa, M., Taketa, M., ... Harada, K. (2001). Construction of a genetic linkage map of the model legume *Lotus japonicus* using an intraspecific F2 population. *DNA Research*, 8(6), 301–310. <https://doi.org/10.1093/dnares/8.6.301>
- Huda, I. N., & Daryono, B. S. (2013). Analisis variasi genetik melon (*Cucumis melo* L.) kultivar gama melon basket dengan metode random amplified polymorphic DNA. *Biogenesis: Jurnal Ilmiah Biologi*, 1(1), 41–50. <https://doi.org/10.24252/bio.v1i1.446>
- Jiang, J., & Gill, B. S. (2006). Current status and the future of fluorescence *in situ* hybridization (FISH) in plant genome research. *Genome*, 49(9), 1057–1068. <https://doi.org/10.1139/G06-076>
- Kato, A., Lamb, J. C., & Birchler, J. A. (2004). Chromosome painting using repetitive DNA sequences as probes for somatic chromosome identification in maize. *Proceedings of the National Academy of Sciences of the United States of America*, 101(37), 13554–13559. <https://doi.org/10.1073/pnas.0403659101>
- Koo, D.-H., Nam, Y.-W., Choi, D., Bang, J.-W., de Jong, H., & Hur, Y. (2010). Molecular cytogenetic mapping of *Cucumis sativus* and *C. melo* using highly repetitive DNA sequences. *Chromosome Research*, 8, 325–336. <https://doi.org/10.1007/s10577-010-9116-0>
- Li, K., Wang, H., Wang, J., Sun, J., Li, Z., & Han, Y. (2016). Divergence between *C. melo* and African *Cucumis* species identified by chromosome painting and rDNA distribution pattern. *Cytogenetic and Genome Research*, 150(2), 150–155. <https://doi.org/10.1159/000453520>

- Agus Budi Setiawan *et al.*: *Cytogenetic Evidences Timun Suri and Cucumber*.....
- Li, Z.-Y., Fu, M.-L., Hu, F.-F., Huang, S.-F., & Song, Y.-C. (2006). Visualization of the ribosomal DNA (45S rDNA) of *Indica* rice with FISH on some phases of cell cycle and extended DNA fibers. *Biocell*, 30(1), 27–32. <https://doi.org/10.32604/biocell.2006.30.027>
- Liu, C., Liu, J., Li, H., Zhang, Z., Han, Y., Huang, S., & Jin, W. (2010). Karyotyping in melon (*Cucumis melo* L.) by cross-species fosmid fluorescence in situ hybridization. *Cytogenetic and Genome Research*, 129(1–3), 241–249. <https://doi.org/10.1159/000314343>
- Lou, Q., Zhang, Y., He, Y., Li, J., Jia, L., Cheng, C., ... Chen, J. (2014). Single-copy gene-based chromosome painting in cucumber and its application for chromosome rearrangement analysis in *Cucumis*. *Plant Journal*, 78(1), 169–179. <https://doi.org/10.1111/tpj.12453>
- Manohar, S. H., & Murthy, H. N. (2012). Estimation of phenotypic divergence in a collection of *Cucumis melo*, including shelf-life of fruit. *Scientia Horticulturae*, 148, 74–82. <https://doi.org/10.1016/j.scienta.2012.09.025>
- Marcon, A. B., Leão Barros, I. C., & Guerra, M. (2005). Variation in chromosome numbers, CMA bands and 45S rDNA sites in species of *Selaginella* (Pteridophyta). *Annals of Botany*, 95(2), 271–276. <https://doi.org/10.1093/aob/mci022>
- Navrátilová, A., Neumann, P., & Macas, J. (2003). Karyotype analysis of four *Vicia* species using *in situ* hybridization with repetitive sequences. *Annals of Botany*, 91(7), 921–926. <https://doi.org/10.1093/aob/mcg099>
- Neves, N., Delgado, M., Silva, M., Caperta, A., Morais-Cecílio, L., & Viegas, W. (2005). Ribosomal DNA heterochromatin in plants. *Cytogenetic and Genome Research*, 109(1–3), 104–111. <https://doi.org/10.1159/000082388>
- Pedrosa-Harand, A., de Almeida, C. C. S., Mosiolek, M., Blair, M. W., Schweizer, D., & Guerra, M. (2006). Extensive ribosomal DNA amplification during Andean common bean (*Phaseolus vulgaris* L.) evolution. *Theoretical and Applied Genetics*, 112, 924–933. <https://doi.org/10.1007/s00122-005-0196-8>
- Penner, G. A., Bush, A., Wise, R., Kim, W., Domier, L., Kasha, K., ... Fedak, G. (1993). Reproducibility of random amplified polymorphic DNA (RAPD) analysis among laboratories. *Genome Research*, 2, 341–345. <https://doi.org/10.1101/gr.2.4.341>
- Pitrat, M., Chauvet, M., & Foury, C. (1999). Diversity, history and production of cultivated cucurbits. *Acta Horticulturae*, 492, 21–28. <https://doi.org/10.17660/ActaHortic.1999.492.1>
- Rahayu, S. E., & Hartana, A. (2002). Biosistematika *Cucumis* (Cucurbitaceae) di Jawa. *Floribunda*, 2(2), 38–43. Retrieved from <http://www.ptti.or.id/journal/index.php/Floribunda/article/view/39>
- Raskina, O., Belyayev, A., & Nevo, E. (2004a). Activity of the En/Spm-like transposons in meiosis as a base for chromosome repatterning in a small, isolated, peripheral population of *Aegilops speltoides* Tausch. *Chromosome Research*, 12(2), 153–161. <https://doi.org/10.1023/B:CHRO.0000013168.61359.43>
- Raskina, O., Belyayev, A., & Nevo, E. (2004b). Quantum speciation in *Aegilops*: Molecular cytogenetic evidence from rDNA cluster variability in natural populations. *Proceedings of the National Academy of Sciences of the United States of America*, 101(41), 14818–14823. <https://doi.org/10.1073/pnas.0405817101>
- Roa, F., & Guerra, M. (2012). Distribution of 45S rDNA sites in chromosomes of plants: Structural and evolutionary implications. *BMC Evolutionary Biology*, 12, 225. <https://doi.org/10.1186/1471-2148-12-225>
- Schwarzacher, T., & Heslop-Harrison, P. (2000). *Practical in situ hybridization*. Oxford: BIOS Scientific Publishers. Retrieved from <https://www.cabdirect.org/cabdirect/abstract/20001610782>
- Sebastian, P., Schaefer, H., Telford, I. R. H., & Renner, S. S. (2010). Cucumber (*Cucumis sativus*) and melon (*C. melo*) have numerous wild relatives in Asia and Australia, and the sister species of melon is from Australia. *Proceedings of the National Academy of Sciences of the United States of America*, 107(32), 14269–14273. <https://doi.org/10.1073/pnas.1005338107>
- Setiawan, A. B. (2018). Molecular cytogenetic studies on satellite DNA and retrotransposon in *Cucumis* species (Thesis). Chiba University.
- Setiawan, A. B., Teo, C. H., Kikuchi, S., Sassa, H., & Koba, T. (2018). An improved method for inducing prometaphase chromosomes in plants. *Molecular Cytogenetics*, 11, 32. <https://doi.org/10.1186/s13039-018-0380-6>
- She, C.-W., Jiang, X.-H., Ou, L.-J., Liu, J., Long, K.-L., Zhang, L.-H., ... Hu, J.-C. (2015). Molecular cytogenetic characterisation and phylogenetic analysis of the seven cultivated *Vigna* species (Fabaceae). *Plant Biology*, 17(1), 268–280.

Agus Budi Setiawan *et al.*: Cytogenetic Evidences Timun Suri and Cucumber.....

<https://doi.org/10.1111/plb.12174>

- Singh, A. K., Kumar, S., Singh, H., Rai, V. P., Singh, B. D., & Pandey, S. (2015). Genetic diversity in Indian snapmelon (*Cucumis melo* var. *momordica*) accessions revealed by ISSR markers. *Plant OMICS*, *8*(1), 9–16. Retrieved from https://www.pomics.com/pandy_8_1_2015_9_16.pdf
- Skroch, P., & Nienhuis, J. (1995). Impact of scoring error and reproducibility RAPD data on RAPD based estimates of genetic distance. *Theoretical and Applied Genetics*, *91*, 1086–1091. <https://doi.org/10.1007/BF00223923>
- Vaio, M., Speranza, P., Valls, J. F., Guerra, M., & Mazzella, C. (2005). Localization of the 5S and 45S rDNA sites and cpDNA sequence analysis in species of the quadrifaria group of *Paspalum* (Poaceae, Paniceae). *Annals of Botany*, *96*(2), 191–200. <https://doi.org/10.1093/aob/mci168>
- Wendel, J. F., Schnabel, A., & Seelanan, T. (1995). Bidirectional interlocus concerted evolution following allopolyploid speciation in cotton (*Gossypium*). *Proceedings of the National Academy of Sciences of the United States of America*, *92*(1), 280–284. <https://doi.org/10.1073/pnas.92.1.280>
- Zhang, Y., Cheng, C., Li, J., Yang, S., Wang, Y., Li, Z., ... Lou, Q. (2015). Chromosomal structures and repetitive sequences divergence in *Cucumis* species revealed by comparative cytogenetic mapping. *BMC Genomics*, *16*, 730. <https://doi.org/10.1186/s12864-015-1877-6>
- Zhang, Z.-T., Yang, S.-Q., Li, Z.-A., Zhang, Y.-X., Wang, Y.-Z., Cheng, C. Y., ... Lou, Q. F. (2016). Comparative chromosomal localization of 45S and 5S rDNAs and implications for genome evolution in *Cucumis*. *Genome*, *59*(7), 449–457. <https://doi.org/10.1139/gen-2015-0207>
- Zhao, X., Lu, J., Zhang, Z., Hu, J., Huang, S., & Jin, W. (2011). Comparison of the distribution of the repetitive DNA sequences in three variants of *Cucumis sativus* reveals their phylogenetic relationships. *Journal of Genetics and Genomics*, *38*(1), 39–45. <https://doi.org/10.1016/j.jcg.2010.12.005>